



cap

## Protocol for the Examination of Specimens From Patients With Neuroendocrine Tumors (Carcinoid Tumors) of the Stomach

**Protocol applies to well-differentiated neuroendocrine tumors of the stomach. Carcinomas with mixed endocrine/glandular differentiation, poorly differentiated carcinomas with neuroendocrine features, and small cell carcinomas are not included.**

---

**Based on AJCC/UICC TNM, 7<sup>th</sup> Edition**

Protocol web posting date: June 2012

### Procedures

- Endoscopic Resection
- Gastrectomy (Partial or Complete)

### Authors

Laura H. Tang, MD, PhD, FCAP\*

Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York, NY

Jordan Berlin, MD

Department of Medicine, Vanderbilt University Medical Center, Nashville, TN

Philip Branton, MD, FCAP

Department of Pathology, Inova Fairfax Hospital, Falls Church, VA

Lawrence J. Burgart, MD, FCAP

Allina Laboratories, Abbott Northwestern Hospital, Minneapolis, MN

David K. Carter, MD, FCAP

Department of Pathology, St. Mary's/Duluth Clinic Health System, Duluth, MN

Carolyn C. Compton, MD, PhD, FCAP

Critical Path Institute, Tucson, AZ

Patrick Fitzgibbons, MD, FCAP

Department of Pathology, St. Jude Medical Center, Fullerton, CA

Wendy L. Frankel, MD, FCAP

Department of Pathology, Ohio State University Medical Center, Columbus, OH

John Jessup, MD

Division of Cancer Treatment and Diagnosis, National Cancer Institute, Bethesda, MD

Sanjay Kakar, MD, FCAP

Department of Pathology, University of California San Francisco and the Veterans Affairs Medical Center, San Francisco, CA

Bruce Minsky, MD

Department of Radiation Oncology, University of Chicago, Chicago, IL

Raouf Nakhleh, MD, FCAP

Department of Pathology, Mayo Clinic, Jacksonville, FL

Kay Washington, MD, PhD, FCAP†

Department of Pathology, Vanderbilt University Medical Center, Nashville, TN

For the Members of the Cancer Committee, College of American Pathologists

\* Denotes primary author. † Denotes senior author. All other contributing authors are listed alphabetically.

© 2012 College of American Pathologists (CAP). All rights reserved.

The College does not permit reproduction of any substantial portion of these protocols without its written authorization. The College hereby authorizes use of these protocols by physicians and other health care providers in reporting on surgical specimens, in teaching, and in carrying out medical research for nonprofit purposes. This authorization does not extend to reproduction or other use of any substantial portion of these protocols for commercial purposes without the written consent of the College.

The CAP also authorizes physicians and other health care practitioners to make modified versions of the Protocols solely for their individual use in reporting on surgical specimens for individual patients, teaching, and carrying out medical research for non-profit purposes.

The CAP further authorizes the following uses by physicians and other health care practitioners, in reporting on surgical specimens for individual patients, in teaching, and in carrying out medical research for non-profit purposes: (1) **Dictation** from the original or modified protocols for the purposes of creating a text-based patient record on paper, or in a word processing document; (2) **Copying** from the original or modified protocols into a text-based patient record on paper, or in a word processing document; (3) The use of a **computerized system** for items (1) and (2), provided that the Protocol data is stored intact as a single text-based document, and is not stored as multiple discrete data fields.

Other than uses (1), (2), and (3) above, the CAP does not authorize any use of the Protocols in electronic medical records systems, pathology informatics systems, cancer registry computer systems, computerized databases, mappings between coding works, or any computerized system without a written license from CAP. Applications for such a license should be addressed to the SNOMED Terminology Solutions division of the CAP.

Any public dissemination of the original or modified Protocols is prohibited without a written license from the CAP.

The College of American Pathologists offers these protocols to assist pathologists in providing clinically useful and relevant information when reporting results of surgical specimen examinations of surgical specimens. The College regards the reporting elements in the "Surgical Pathology Cancer Case Summary" portion of the protocols as essential elements of the pathology report. However, the manner in which these elements are reported is at the discretion of each specific pathologist, taking into account clinician preferences, institutional policies, and individual practice.

The College developed these protocols as an educational tool to assist pathologists in the useful reporting of relevant information. It did not issue the protocols for use in litigation, reimbursement, or other contexts. Nevertheless, the College recognizes that the protocols might be used by hospitals, attorneys, payers, and others. Indeed, effective January 1, 2004, the Commission on Cancer of the American College of Surgeons mandated the use of the required data elements of the protocols as part of its Cancer Program Standards for Approved Cancer Programs. Therefore, it becomes even more important for pathologists to familiarize themselves with these documents. At the same time, the College cautions that use of the protocols other than for their intended educational purpose may involve additional considerations that are beyond the scope of this document.

The inclusion of a product name or service in a CAP publication should not be construed as an endorsement of such product or service, nor is failure to include the name of a product or service to be construed as disapproval.

## CAP Stomach NET Protocol Revision History

---

### Version Code

The definition of the version code can be found at [www.cap.org/cancerprotocols](http://www.cap.org/cancerprotocols).

**Version:** StomachNET 3.2.0.0

### Summary of Changes

The following changes have been made since the November 2011 release.

### Endoscopic Resection, Gastrectomy

#### **Histologic Type; Alternate Histologic Classification; Histologic Grade**

These three reporting elements were combined into one, and the word "checklist" was changed to "protocol" as follows:

#### **Histologic Type and Grade<sup>#</sup>**

- Not applicable
- Well-differentiated neuroendocrine tumor; GX: Grade cannot be assessed
- Well-differentiated neuroendocrine tumor; G1: Low grade (carcinoid)
- Well-differentiated neuroendocrine tumor; G2: Intermediate grade (atypical carcinoid)
- Other (specify): \_\_\_\_\_

<sup>#</sup> For poorly differentiated (high-grade) neuroendocrine carcinomas (G3), the College of American Pathologists (CAP) protocol for carcinoma of the stomach<sup>1</sup> should be used.

#### **Margins**

"Lateral Mucosal Margins" was changed to "Mucosal Margins." Deep Margin and Mucosal Margins: added "required only if applicable," changed "Uninvolved/Involved by tumor" to "Uninvolved/Involved by neuroendocrine tumor," and deleted "Not applicable" data element. Other Margins: added "required only if applicable" and deleted "Not applicable" data element, as follows:

Deep Margin (a endoscopic resections) (required only if applicable)

- Cannot be assessed
- Uninvolved by neuroendocrine tumor
- Involved by neuroendocrine tumor

Mucosal Margins (endoscopic resections) (required only if applicable)

- Uninvolved by neuroendocrine tumor
- Involved by neuroendocrine tumor

Other Margin(s) (required only if applicable)

Specify margin(s): \_\_\_\_\_

- Cannot be assessed
- Uninvolved by neuroendocrine tumor
- Involved by neuroendocrine tumor
- +  Involved by neuroendocrine cell hyperplasia/dysplasia

#### **Ancillary Studies**

Reporting on Ki-67 was updated, changing >2% to 3% and adding "specify," as follows:

- +  Ki-67 labeling index (specify: \_\_\_\_\_)
  - +  ≤2%
  - +  3% to 20%
  - +  >20%

### **Explanatory Notes**

"Neoplasm" was changed to "tumor."

### **D. Histologic Type**

The first sentence was changed to the following:

The World Health Organization (WHO) classifies neuroendocrine neoplasms as well-differentiated neuroendocrine tumors (either the primary tumor or metastasis) and poorly differentiated neuroendocrine carcinomas.<sup>5-8</sup>

### **Alternative Classification Based Upon WHO Classification: Neuroendocrine Tumors of the Appendix**

This section was deleted.

### **E. Histologic Grade**

"Mitotic count" was changed to "mitotic rate."

Ki-67 Index (%): >2 was changed to 3.

The last sentence of the first paragraph was changed to:

The following grading system is recommended by both the European Neuroendocrine Tumor Society (ENETS) and the WHO<sup>8,9</sup>:

### **References**

Reference #8 was updated.

**Surgical Pathology Cancer Case Summary**

---

Protocol web posting date: June 2012

**STOMACH: Endoscopic Resection, Gastrectomy (Note A)****Select a single response unless otherwise indicated.****Specimen (select all that apply)**

- Stomach  
 Portion of stomach  
      Gastric body  
      Gastric antrum  
      Not specified  
 Distal esophagus  
 Proximal duodenum  
 Other (specify): \_\_\_\_\_  
 Not specified

**Procedure**

- Endoscopic resection  
 Partial gastrectomy, proximal  
 Partial gastrectomy, distal  
 Partial gastrectomy, other (specify): \_\_\_\_\_  
 Total gastrectomy  
 Other (specify): \_\_\_\_\_  
 Not specified

**+ Specimen Size (if applicable)**

+ Specify: \_\_\_ (length) x \_\_\_ x \_\_\_ cm

**Tumor Site (select all that apply) (Note B)**

- Gastric cardia  
 Gastric fundus  
 Gastric body  
 Gastric antrum  
 Other (specify): \_\_\_\_\_  
 Not specified

**Tumor Size (Note C)**

Greatest dimension: \_\_\_ cm (specify size of largest tumor if multiple tumors are present)

+ Additional dimensions: \_\_\_ x \_\_\_ cm

 Cannot be determined (see "Comment")**Tumor Focality**

- Unifocal  
 Multifocal (specify number of tumors: \_\_\_\_\_)  
 Cannot be determined

**Histologic Type and Grade (Notes D and E)#**

- Not applicable  
 Well-differentiated neuroendocrine tumor; GX: Grade cannot be assessed  
 Well-differentiated neuroendocrine tumor; G1: Low grade (carcinoid)  
 Well-differentiated neuroendocrine tumor; G2: Intermediate grade (atypical carcinoid)  
 Other (specify): \_\_\_\_\_

# For poorly differentiated (high-grade) neuroendocrine carcinomas (G3), the College of American Pathologists (CAP) protocol for carcinoma of the stomach<sup>1</sup> should be used.

**Mitotic Rate (Note E)**

- Specify: \_\_\_/10 high-power fields (HPF)  
 Cannot be determined

**Microscopic Tumor Extension**

- Cannot be assessed  
 No evidence of primary tumor  
 Tumor invades lamina propria  
 Tumor invades into but not through muscularis mucosae  
 Tumor invades submucosa  
 Tumor invades muscularis propria  
 Tumor invades subserosal tissue without involvement of visceral peritoneum  
 Tumor penetrates serosa (visceral peritoneum)  
 Tumor directly invades adjacent structures (specify: \_\_\_\_\_)  
 Tumor penetrates to the surface of the visceral peritoneum (serosa) and directly invades adjacent structures (specify: \_\_\_\_\_)

**Margins (select all that apply)**

- If all margins uninvolved by neuroendocrine tumor:  
 Distance of tumor from closest margin: \_\_\_ mm or \_\_\_ cm  
 Specify margin: \_\_\_\_\_

Proximal Margin

- Cannot be assessed  
 Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor  
 +  Involved by neuroendocrine cell hyperplasia/dysplasia

Distal Margin

- Cannot be assessed  
 Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor  
 +  Involved by neuroendocrine cell hyperplasia/dysplasia

Omental (Radial) Margin (Note F)

- Cannot be assessed  
 Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor

+ Data elements preceded by this symbol are not required. However, these elements may be clinically important but are not yet validated or regularly used in patient management.

Deep Margin (endoscopic resections) (required only if applicable)

- Cannot be assessed  
 Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor

Mucosal Margins (endoscopic resections) (required only if applicable)

- Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor

Other Margin(s) (required only if applicable)

Specify margin(s): \_\_\_\_\_

- Cannot be assessed  
 Uninvolved by neuroendocrine tumor  
 Involved by neuroendocrine tumor  
+  Involved by neuroendocrine cell hyperplasia/dysplasia

#### **Lymph-Vascular Invasion**

- Not identified  
 Present  
 Indeterminate

#### **+ Perineural Invasion**

- +  Not identified  
+  Present  
+  Indeterminate

#### **Pathologic Staging (pTNM) (Note G)**

TNM Descriptors (required only if applicable) (select all that apply)

- m (multiple primary tumors)  
 r (recurrent)  
 y (posttreatment)

Primary Tumor (pT)

- pTX: Primary tumor cannot be assessed  
 pT0: No evidence of primary tumor  
 pTis: Carcinoma in situ/dysplasia (tumor size less than 0.5 mm), confined to mucosa  
 pT1: Tumor invades lamina propria or submucosa and 1 cm or less in size  
 pT2: Tumor invades muscularis propria or more than 1 cm in size  
 pT3: Tumor penetrates subserosa  
 pT4: Tumor invades visceral peritoneum (serosal) or other organs or adjacent structures

Regional Lymph Nodes (pN)

- Cannot be assessed  
 pN0: No regional lymph node metastasis  
 pN1: Metastasis in regional lymph nodes  
 No nodes submitted or found

+ Data elements preceded by this symbol are not required. However, these elements may be clinically important but are not yet validated or regularly used in patient management.

*Number of Lymph Nodes Examined*

Specify: \_\_\_\_

\_\_\_ Number cannot be determined (explain): \_\_\_\_\_

*Number of Lymph Nodes Involved*

Specify: \_\_\_\_

\_\_\_ Number cannot be determined (explain): \_\_\_\_\_

Distant Metastasis (pM)

\_\_\_ Not applicable

\_\_\_ pM1: Distant metastasis

+ Specify site(s), if known: \_\_\_\_\_

**+ Ancillary Studies (select all that apply) (Notes E and H)**

+ \_\_\_ Ki-67 labeling index (specify: \_\_\_\_)

+ \_\_\_ ≤2%

+ \_\_\_ 3% to 20%

+ \_\_\_ &gt;20%

+ \_\_\_ Other (specify): \_\_\_\_\_

+ \_\_\_ Not performed

**+ Additional Pathologic Findings (select all that apply) (Note I)**

+ \_\_\_ Atrophic gastritis

+ \_\_\_ Intestinal metaplasia of gastric mucosa

+ \_\_\_ Glandular dysplasia of gastric mucosa

+ \_\_\_ Endocrine cell hyperplasia

+ \_\_\_ Absence of parietal cells

+ \_\_\_ Tumor necrosis

+ \_\_\_ Other, specify: \_\_\_\_\_

**+ Comment(s)**



## Explanatory Notes

### A. Application and Tumor Location

This protocol applies to well-differentiated neuroendocrine tumors (carcinoid tumors) of the stomach. Poorly differentiated neuroendocrine carcinomas, small cell carcinomas, and tumors with mixed glandular/neuroendocrine differentiation are not included.

Because of site-specific similarities in histology, immunohistochemistry, and histochemistry, neuroendocrine tumors of the digestive tract have traditionally been subdivided into those of foregut, midgut, and hindgut origin (Table 1). In general, the distribution pattern along the gastrointestinal (GI) tract parallels that of the progenitor cell type, and the anatomic site of origin of GI neuroendocrine tumors is an important predictor of clinical behavior.<sup>2</sup>

**Table 1. Site of Origin of Gastrointestinal Neuroendocrine Tumors**

	Foregut Tumors	Midgut Tumors	Hindgut Tumors
<i>Site</i>	<i>Stomach, Proximal Duodenum</i>	<i>Jejunum, Ileum, Appendix, Proximal Colon</i>	<i>Distal Colon, Rectum</i>
Immunohistochemistry			
Chromogranin A	86%-100% +	82%-92% +	40%-58% +
Neuron-Specific Enolase (NSE)	90%-100% +	95%-100% +	80%-87% +
Synaptophysin	50% +	95%-100% +	94%-100% +
Serotonin	33% + <sup>11,12</sup>	86% + <sup>11,12</sup>	45%-83% + <sup>3-5,12</sup>
Other Immunohistochemical Markers	Rarely, + for pancreatic polypeptide, histamine, gastrin, somatostatin, vasoactive intestinal peptide (VIP), or adrenocorticotrophic hormone (ACTH)	Prostatic acid phosphatase + in 20%-40% <sup>11,12</sup>	Prostatic acid phosphatase + in 20%-82% <sup>3-5,12</sup>
Carcinoid Syndrome	Rare	5%-39% <sup>6,7</sup>	Rare

### B. Site-Specific Features

Gastric neuroendocrine tumors are divided into 4 types.<sup>3</sup> Type 1 tumors arising in the setting of atrophic gastritis with associated hypergastrinemia are the most common. These lesions are composed of enterochromaffin-like (ECL) cells and are usually found as multiple small nodules in the body of the stomach and limited to mucosa and submucosa. Type 1 lesions are generally benign and may regress following antrectomy; lymph node metastases are very rare and occur only when the tumors are large (greater than 2 cm) and infiltrate the muscularis propria.

Type 2 gastric neuroendocrine tumors are rare. These multifocal small tumors, which are associated with multiple endocrine neoplasia (MEN) type 1 with Zollinger-Ellison syndrome, develop in the body of the stomach, are usually smaller than 1.5 cm, and are confined to the mucosa or submucosa. However, in contrast to type 1 tumors, 30% metastasize. Tumors greater than 2 cm and invading the muscularis propria and exhibiting vascular invasion are more likely to metastasize.

Type 3 gastric neuroendocrine tumors, the second most common neuroendocrine tumor in the stomach, are sporadic solitary tumors that are unassociated with atrophic gastritis or endocrine cell hyperplasia. These tumors may occur anywhere in the stomach. Metastasis is associated with larger mean size, angioinvasion, and invasion of muscularis propria. Surgical resection is usually advised for

solitary gastric carcinoid tumors, particularly those larger than 2.0 cm, but tumors smaller than 1.0 cm have been rarely reported to metastasize.<sup>4</sup>

Type 4 gastric neuroendocrine tumors are rare high-grade neuroendocrine carcinomas that are usually bulky tumors with metastases at diagnosis (the CAP cancer protocol for gastric carcinoma applies<sup>1</sup>).

### C. Tumor Size

For neuroendocrine tumors in any part of the gastrointestinal tract, size greater than 2.0 cm is associated with a higher risk of lymph node metastasis. In the stomach, types 3 and 4 neuroendocrine tumors are significantly larger than type 1 tumors,<sup>3</sup> which usually measure 1 cm or less<sup>5,6</sup> (Table 2). Tumor size correlates with depth of invasion for gastric neuroendocrine tumors, with larger tumors more likely to be deeply infiltrative and thus at higher risk for metastases. Nodules measuring 0.5 mm or larger are defined as neuroendocrine tumors; lesions measuring less than 0.5 mm are regarded as representing in situ tumor, neuroendocrine cell dysplasia, or hyperplasia.

**Table 2. Types of Gastric Neuroendocrine Tumors**

	<b>Type 1</b>	<b>Type 2</b>	<b>Type 3</b>	<b>Type 4</b>
Frequency	70%-80% of cases	Rare	10%-15% of cases	Rare
Multiplicity	Multifocal	Multifocal	Solitary	Solitary
Size	0.5-1.0 cm	~1.5 cm or less	Variable; one-third are larger than 2 cm	Large
Location	Corpus	Corpus	Anywhere in stomach	Anywhere in stomach
Associations	Hypergastrinemic states; chronic atrophic gastritis, enterochromaffin-like (ECL) cell hyperplasia, pernicious anemia	Multiple endocrine neoplasia (MEN) type 1, with hypergastrinemia or Zollinger-Ellison syndrome	Sporadic	Sporadic
Clinical Behavior	Usually benign	30% metastasize	71% of tumors >2 cm with muscularis propria and vascular invasion have lymph node metastases	High-grade carcinoma. Metastases common; poor prognosis
Demographic Profile	70%-80% are females in their 50s and 60s	Equally in males and females, mean age 50 y	More common in males, mean age 55 y	More common in males

### D. Histologic Type

The World Health Organization (WHO) classifies neuroendocrine neoplasms as well-differentiated neuroendocrine tumors (either the primary tumor or metastasis) and poorly differentiated neuroendocrine carcinomas.<sup>5-8</sup> Historically, well-differentiated neuroendocrine tumors have been referred to as carcinoid tumors, a term which may cause confusion because clinically a carcinoid tumor is a serotonin-producing tumor associated with functional manifestations of carcinoid syndrome.

Classification of neuroendocrine tumors (NETs) is based upon size, functionality, site, and invasion. Functioning tumors are those associated with clinical manifestations of hormone production or secretion of measurable amounts of active hormone; immunohistochemical demonstration of hormone production is not equivalent to clinically apparent functionality.

### Histologic Patterns

Although specific histologic patterns in well-differentiated neuroendocrine tumors, such as trabecular, insular, and glandular, roughly correlate with tumor location, these patterns have not been clearly shown independently to predict response to therapy or risk of nodal metastasis and are rarely reported in clinical practice.

### E. Histologic Grade

Cytologic atypia in low-grade neuroendocrine tumors has no impact on clinical behavior of these tumors. However, grading systems based on mitotic activity have been shown to have utility for foregut tumors. The following grading system is recommended by both the European Neuroendocrine Tumor Society (ENETS) and the WHO<sup>8,9</sup>:

Grade	Mitotic Rate (per 10 HPF) #	Ki-67 Index (%)##
G1	<2	≤2
G2	2 to 20	3 to 20
G3	>20	>20

# Mitotic rate should be based upon counting 50 high-power (40x objective) fields in the area of highest mitotic activity and reported as number of mitoses per 10 HPF.

## Ki-67 index is reported as percentage of positive tumor cells in area of highest nuclear labeling. It has been recommended that 2000 tumor cells be counted to determine the Ki-67 index<sup>9</sup>; however, this practice may not be practical for routine clinical purposes, and it is acceptable to estimate the labeling index.

G1 and G2 are well-differentiated tumors with diffuse intense chromogranin/synaptophysin positivity. Punctate necrosis is more typical of G2 tumors. G3 tumors are high-grade neuroendocrine carcinomas (the CAP carcinoma protocol for carcinomas of the stomach applies<sup>1</sup>).

### F. Circumferential (Radial) Margin

For surgical resection specimens, margins include the proximal, distal, and radial margins. The radial margins represent the nonperitonealized soft tissue margins closest to the deepest penetration of tumor. In the stomach, the lesser omental (hepatoduodenal and hepatogastric ligaments) and greater omental resection margins are the only radial margins. For endoscopic resection specimens, margins include mucosal margins and the deep margin of resection. It may be helpful to mark the margin(s) closest to the tumor with ink. Margins marked by ink should be designated in the macroscopic description.

### G. TNM and Anatomic Stage/Prognostic Groupings

The TNM staging system for gastric neuroendocrine tumors of the American Joint Committee on Cancer (AJCC) and the International Union Against Cancer (UICC) is recommended.<sup>10</sup>

By AJCC/UICC convention, the designation "T" refers to a primary tumor that has not been previously treated. The symbol "p" refers to the pathologic classification of the TNM, as opposed to the clinical classification, and is based on gross and microscopic examination. pT entails a resection of the primary tumor or biopsy adequate to evaluate the highest pT category, pN entails removal of nodes adequate to validate lymph node metastasis, and pM implies microscopic examination of distant lesions. Clinical classification (cTNM) is usually carried out by the referring physician before treatment during initial evaluation of the patient or when pathologic classification is not possible.

Pathologic staging is usually performed after surgical resection of the primary tumor. Pathologic staging depends on pathologic documentation of the anatomic extent of disease, whether or not the primary tumor has been completely removed. If a biopsied tumor is not resected for any reason (eg, when

technically unfeasible) and if the highest T and N categories or the M1 category of the tumor can be confirmed microscopically, the criteria for pathologic classification and staging have been satisfied without total removal of the primary cancer.

### TNM Descriptors

For identification of special cases of TNM or pTNM classifications, the “m” suffix and “y,” “r,” and “a” prefixes are used. Although they do not affect the stage grouping, they indicate cases needing separate analysis.

The “m” suffix indicates the presence of multiple primary tumors in a single site and is recorded in parentheses: pT(m)NM.

The “y” prefix indicates those cases in which classification is performed during or following initial multimodality therapy (ie, neoadjuvant chemotherapy, radiation therapy, or both chemotherapy and radiation therapy). The cTNM or pTNM category is identified by a “y” prefix. The ycTNM or ypTNM categorizes the extent of tumor actually present at the time of that examination. The “y” categorization is not an estimate of tumor prior to multimodality therapy (ie, before initiation of neoadjuvant therapy).

The “r” prefix indicates a recurrent tumor when staged after a documented disease-free interval, and is identified by the “r” prefix: rTNM.

The “a” prefix designates the stage determined at autopsy: aTNM.

### N Category Considerations

The specific nodal areas of the stomach are listed below.<sup>10</sup>

Greater curvature of stomach: Greater curvature, greater omental, gastroduodenal, gastroepiploic, pyloric, and pancreaticoduodenal

Pancreatic and splenic area: Pancreaticolienal, peripancreatic, splenic

Lesser curvature of stomach: Lesser curvature, lesser omental, left gastric, cardioesophageal, common hepatic, celiac, and hepatoduodenal

Involvement of other intra-abdominal lymph nodes, such as hepatoduodenal, retropancreatic, mesenteric, and para-aortic, is classified as distant metastasis.<sup>10</sup>

### TNM Anatomic Stage/Prognostic Groupings

Stage 0	Tis	N0	M0 <sup>#</sup>
Stage I	T1	N0	M0
Stage IIa	T2	N0	M0
Stage IIb	T3	N0	M0
Stage IIIa	T4	N0	M0
Stage IIIb	Any T	N1	M0
Stage IV	Any T	Any N	M1

<sup>#</sup> M0 is defined as no distant metastasis.

### H. Ancillary Studies

Immunohistochemistry and other ancillary techniques are generally not required to diagnose well-differentiated neuroendocrine tumors. Specific markers that may be used to establish neuroendocrine differentiation include chromogranin A, neuron-specific enolase, synaptophysin, and CD56.<sup>6</sup> Because of their relative sensitivity and specificity, chromogranin A and synaptophysin are recommended.

Immunohistochemistry for Ki-67 may be useful in establishing tumor grade (Note E) and prognosis<sup>8</sup> but is not currently considered standard of care.<sup>6</sup>

Immunohistochemistry for specific hormone products, such as glucagon, gastrin, and somatostatin, may be of interest in some cases. However, immunohistochemical demonstration of hormone production does not equate with clinical functionality of the tumor.

### I. Additional Pathologic Findings

Most gastric neuroendocrine tumors arise in the setting of chronic atrophic gastritis (see Note B). Atrophic gastritis may be associated with glandular dysplasia, and in rare cases, gastric adenocarcinoma. Coagulative tumor necrosis, usually punctate, may indicate more aggressive behavior<sup>9</sup> and should be reported.

### References

1. Washington K, Berlin J, Branton P, et al. Protocol for the examination of specimens from patients with carcinoma of the stomach. In: *Reporting on Cancer Specimens: Case Summaries and Background Documentation*. Northfield, IL: College of American Pathologists; 2009.
2. Rorstad O. Prognostic indicators for carcinoid neuroendocrine tumors of the gastrointestinal tract. *J Surg Oncol*. Mar 1 2005;89(3):151-160.
3. Borch K, Ahren B, Ahlman H, Falkmer S, Granerus G, Grimelius L. Gastric carcinoids: biologic behavior and prognosis after differentiated treatment in relation to type. *Ann Surg*. 2005;242(1):64-73.
4. Xie SD, Wang LB, Song XY, Pan T. Minute gastric carcinoid with regional lymph node metastasis: a case report and review of the literature. *World J Gastroenterol*. 2004;10(16):2461-2463.
5. Graeme-Cook F. Neuroendocrine tumors of the GI tract and appendix. In: Odze RD, Goldblum JR, Crawford JM, eds. *Surgical Pathology of the GI Tract, Liver, Biliary Tract, and Pancreas*. Philadelphia, PA: WB Saunders; 2004: 483-504.
6. Williams GT. Endocrine tumours of the gastrointestinal tract: selected topics. *Histopathology*. 2007;50(1):30-41.
7. Kloppel G, Perren A, Heitz PU. The gastroenteropancreatic neuroendocrine cell system and its tumors: the WHO classification. *Ann N Y Acad Sci*. 2004;1014:13-27.
8. Bosman FT, Carneiro F, Hruban RH, Theise ND, eds. *WHO Classification of Tumours of the Digestive System*. Geneva, Switzerland: WHO Press; 2010.
9. Rindi G, Kloppel G, Ahlman H, et al; and all other Frascati Consensus Conference participants; European Neuroendocrine Tumor Society (ENETS). TNM staging of foregut (neuro)endocrine tumors: a consensus proposal including a grading system. *Virchows Arch*. 2006;449(4):395-401.
10. Edge SB, Byrd DR, Carducci MA, Compton CC. *AJCC Cancer Staging Manual*. 7th ed. New York, NY: Springer; 2009.
11. Kimura N, Sasano N. Prostate-specific acid phosphatase in carcinoid tumors. *Virchows Arch A Pathol Anat Histopathol*. 1986;410(3):247-251.
12. Nash SV, Said JW. Gastroenteropancreatic neuroendocrine tumors: a histochemical and immunohistochemical study of epithelial (keratin proteins, carcinoembryonic antigen) and neuroendocrine (neuron-specific enolase, bombesin and chromogranin) markers in foregut, midgut, and hindgut tumors. *Am J Clin Pathol*. 1986;86(2):415-422.